

 Please cut along the dotted lines before submitting to a pharmacy.

Date Required: \_\_\_\_\_ Ship To:  Patient  MD Office  Other: \_\_\_\_\_

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 City, State, Zip: \_\_\_\_\_  
 Home Phone: \_\_\_\_\_  
 Cell Phone: \_\_\_\_\_  
 Date of Birth: \_\_\_\_\_ Gender: \_\_\_\_\_  
 Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**PRESCRIBER INFORMATION**

Prescriber Name: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 City, State, Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_  
 Fax: \_\_\_\_\_  
 DEA #: \_\_\_\_\_ NPI #: \_\_\_\_\_  
 Contact Person: \_\_\_\_\_

**INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card.)**

Primary Insurance: \_\_\_\_\_ ID: \_\_\_\_\_ Group: \_\_\_\_\_  
 Secondary Insurance: \_\_\_\_\_ ID: \_\_\_\_\_ Group: \_\_\_\_\_  
 Prescription Card: \_\_\_\_\_ ID: \_\_\_\_\_ BIN: \_\_\_\_\_ PCN: \_\_\_\_\_

**To better serve your patient and facilitate insurance authorization, please complete the pertinent sections:**

**PATIENT DIAGNOSIS/CLINICAL INFORMATION**

M06.9 Rheumatoid Arthritis  
 M45.9 Ankylosing Spondylitis  
 M32.9 Systemic Lupus Erythematosus  
 M08.00 Unspecified Juvenile Rheumatoid Arthritis  
 L40.0 Moderate to Severe Plaque Psoriasis  
 L40.50 Psoriatic Arthritis  
 L40.59 Psoriasis with Arthropathy  
 Other: \_\_\_\_\_

TB/PPD Test:  Positive  Negative  Date Read: \_\_\_\_\_  
 Weight: \_\_\_\_\_  kg  lbs Height: \_\_\_\_\_  cm  in %BSA: \_\_\_\_\_  
 Allergies: \_\_\_\_\_  NKDA  
 Pharmacy to coordinate home health nursing visit as necessary:  Yes  No  
 Home health nursing coordination not necessary:  
 Reason:  MD office to administer to patient  
 Home health nursing already coordinated

Prior Medication Failed: \_\_\_\_\_ Length of Treatment: \_\_\_\_\_ Reason for Discontinuation: \_\_\_\_\_

**PRESCRIPTION INFORMATION**

Medication:	Dose/Strength:	Directions:	Quantity:	Refills:
<input type="checkbox"/> Actemra®	<input type="checkbox"/> 80 mg per 4mL <input type="checkbox"/> 200 mg per 10mL <input type="checkbox"/> 400 mg per 20mL	<input type="checkbox"/> Induction Dose: Infuse 4 mg per kg every 4 weeks <input type="checkbox"/> Maintenance Dose: Infuse up to 8 mg per kg every 4 weeks based on clinical response <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Benlysta®	<input type="checkbox"/> 120mg per vial <input type="checkbox"/> 400mg per vial	<input type="checkbox"/> Induction Dose: 10mg per kg. Dose = _____ mg at 2 week intervals for the first 3 doses and at 4-week intervals thereafter. Infuse IV over 1 hour <input type="checkbox"/> Maintenance Dose: 10mg per kg. Dose = _____ mg every 4 weeks. Infuse IV over 1 hour		
<input type="checkbox"/> Orencia®	<input type="checkbox"/> 250mg vial	<input type="checkbox"/> Infuse _____ mg at weeks 0, 2 and 4, then every 4 weeks thereafter <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Inflectra®	<input type="checkbox"/> 100 mg vial	<input type="checkbox"/> INITIAL: Infuse IV _____ mg per kg (Dose _____ mg) at 0, 2, and 6 weeks (Quantity: _____) <input type="checkbox"/> MAINTENANCE: Infuse IV _____ mg per kg (Dose _____ mg) every _____ weeks (Quantity: _____) <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Remicade®				
<input type="checkbox"/> Renflexis™		<input type="checkbox"/> Pharmacist will round to the nearest 100 <input type="checkbox"/> Give exact dose (do NOT round)		
<input type="checkbox"/> Rituxan®	<input type="checkbox"/> 100mg vial <input type="checkbox"/> 500mg vial	<input type="checkbox"/> Infuse two doses of 1000 mg separated by 2 weeks <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Simponi Aria®	<input type="checkbox"/> 50mg per 4mL	<input type="checkbox"/> Infuse 2 mg per kg over 30 minutes at weeks 0 and 4, then every 8 weeks thereafter <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Other:				

**Premedications & Other Medications:**

▶ Infusion supplies as per protocol  
 ▶ Anaphylaxis Kit as per protocol

Acetaminophen: \_\_\_\_\_ mg PO prior to infusion  
 Diphenhydramine: \_\_\_\_\_ mg  
 250mL 0.9% NaCl for hydration  PO  IV  
 Other: \_\_\_\_\_

**Flush Protocol:**

▶ NaCl 0.9% 10mL  
 ▶ Before and after infusion

*By signing this form and using this pharmacy's services, you are authorizing this pharmacy to serve as your prior authorization designated agent in dealing with prescription and medical insurance companies.*

**Prescriber Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_