

 Please cut along the dotted lines before submitting to a pharmacy.

Date Required: _____ Ship To: Home Office Other: _____

PATIENT INFORMATION

Patient Name: _____
 Address: _____
 City, State, Zip: _____
 Home Phone: _____
 Cell Phone: _____
 Date of Birth: _____ Gender: _____
 Emergency Contact: _____ Phone: _____

PRESCRIBER INFORMATION

Prescriber Name: _____
 Address: _____
 City, State, Zip: _____
 Phone: _____
 Fax: _____
 DEA #: _____ NPI #: _____
 Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card.)

Primary Insurance: _____ ID: _____ Group: _____
 Secondary Insurance: _____ ID: _____ Group: _____
 Prescription Card: _____ ID: _____ BIN: _____ PCN: _____

To better serve your patient and facilitate insurance authorization, please complete the pertinent sections:

DIAGNOSIS

E05.00 Thyrotoxicosis with diffuse goiter without thyrotoxic crisis or storm (hyperthyroidism)
 Other: _____

MEDICAL HISTORY

Patient Weight: _____ kg lbs Height: _____ cm in
 Allergies: _____
 Line Access: Peripheral Port
 Delivery Method: Infusion Pump Other: _____
 Therapy Start Date: _____ Therapy End Date: _____
 Nursing Coordination:
 Pharmacy to coordinate home health nursing visit as necessary: Yes No
 Home health nursing coordination not necessary. Reason:
 MD office to administer to patient
 Home health nursing already coordinated
 Lab Orders: _____

PRESCRIPTION INFORMATION

Tepezza® (Teprotumumab-trbw) Prescription:	Quantity:	Refills:
Initiate services beginning with Dose No. _____ as indicated below:		
<input type="checkbox"/> Dose 1: Infuse 10 mg/kg IV over 90 minutes, then 3 weeks later...		
<input type="checkbox"/> Dose 2: Infuse 20 mg/kg IV over 90 minutes, then 3 weeks later...		
<input type="checkbox"/> Dose 3 through 8: Infuse 20 mg/kg IV over 60 to 90 minutes (as tolerated by patient) every 3 weeks x 6 doses.		

PREMEDICATION ORDERS/OTHER MEDICATIONS

<p>Flush Protocol</p> <p>Peripheral:</p> <p><input type="checkbox"/> NaCl 0.9% 5mL <input type="checkbox"/> NaCl 0.9% 10mL</p> <p>Premedications & Other Medications</p> <p><input type="checkbox"/> Infusion supplies as per protocol <input type="checkbox"/> Anaphylaxis Kit orders as per protocol</p>	<p>Implanted Port:</p> <p><input type="checkbox"/> NaCl 0.9% 5 to 10mL pre-/post-use and 10 to 20mL pre-/post-lab draw <input type="checkbox"/> Other: _____ <input type="checkbox"/> Heparin (100 unit/mL) 3 to 5 mL post-use <input type="checkbox"/> For maintenance, heparin (100 unit/mL) 3 to 5mL every 24 hr if accessed or weekly to monthly if not accessed <input type="checkbox"/> Acetaminophen 650 mg PO prior to infusion <input type="checkbox"/> Diphenhydramine 25 mg PO</p>
--	--

ADDITIONAL COMMENTS:

Prescriber Signature: _____ **Date:** _____