## Vyvgart<sup>®</sup> & Vyvgart<sup>®</sup> Hytrulo Enrollment Form

$\gg$ Please cut along the dotted lines before submitting to a pha	armacy.		77-828-3940
Date Required:	Ship To: Patient	MD Office Other:	
PATIENT INFORMATION		PRESCRIBER INFORMATION	
Patient Name:		Prescriber Name:	
Address:		Address:	
City, State, Zip:		City, State, Zip:	
Home Phone:		_ Phone:	
Cell Phone:		Fax: DEA #: NPI #:	
Date of Birth: Gender:		DEA #: INTI # Contact Person:	
Emergency Contact: Phone		ing and a second s	
		front and back of insurance and prescription drug card.)	
Primary Insurance:			
Secondary Insurance: Prescription Card:		ID:         Group:           BIN:         PCN:	
	ID	BIN: PCN:	
To better serve your patient and	facilitate insuranc	e authorization, please complete the pertinent sections:	
DIAGNOSIS		MEDICAL HISTORY	
G70.00 Myasthenia Gravis without (acute) exacerb	ation	Patient Weight: kg lbs Height: cm	in
G70.01 Myasthenia Gravis with (acute) exacerbati		Allergies: Kg ibs fielgitt eff	111
-		Line Access: Peripheral PICC Port	
Clinical/Progress notes with supporting diagnosis		Delivery Method: Infusion Pump Other:	
H&P		Therapy End Date: Therapy End Date:	
Positive serologic test for anti-AChR antibody for gl		Nursing Coordination:	
MG-ADL Score:		<ul> <li>Pharmacy to coordinate home health</li> </ul>	
MGFA classification: Patient's demographics, including insurance inform	ation	<ul> <li>nursing visit as necessary: Yes No</li> </ul>	
Please attach original prescription orders	ation	Home health nursing coordination not necessary. Reason:	
Current medications:		MD office to administer to patient	
Previous therapies: eculizumab rituximab	IVIG	<ul> <li>Home health nursing already coordinated</li> </ul>	
oral corticosteroids non-steroidal ISTs	IVIG		
Previous live vaccine:	Data		
	_ Date:	-	
	PRESCRIPTIC	ON INFORMATION	
Vyvgart® (efgartigimod alfa-fcab) Prescription:			Dispense:
10mg/kg body weight weekly for 4 doses. (Once weekly x 4 weeks.) Maximum dose: 1.2gm. Subsequent treatment cycles may be administered based on clinical evaluation and no sooner than 50 days from the start of the previous treatment cycle.			4 doses
Subsequent treatment cycles to be determined			no refills
Subsequent treatment cycles to be every 50 days from the start of the previous treatment cycle			7 refills
Subsequent treatment cycles to be every days from the start of the treatment cycle			refills
Vyvgart® Hytrulo (efgartigimod alfa and hyaluronidase-qvfc) Prescription:			
1,008 mg/11,200 units (5.6mL) subcutaneous injection over 30-90 seconds once weekly for 4 weeks. Subsequent treatment cycles based on clinical evaluation; no sooner than 50 days from the start of the previous treatment cycle. To be administered by a healthcare professional with 30 minutes post observation period.			4 doses
Subsequent treatment cycles to be determined			no refills
Subsequent treatment cycles to be every 50 days from the start of the previous treatment cycle			7 refills
Subsequent treatment cycles to be every	days from the start of th	e treatment cycle	refills
	ORDERS/OTH	IER MEDICATIONS	
<ul> <li>Orders are initiated:</li> </ul>			
<ul> <li>Infusion supplies as per protocol</li> <li>A perbularia Kit ordera as per protocol</li> </ul>	Flush Protocol (Fe ► NaCl 0.9% 10	or Vyvgart® intravenous infusion only)	
<ul> <li>Anaphylaxis Kit orders as per protocol</li> </ul>	<ul> <li>NaCl 0.9% 101</li> <li>Before and after</li> </ul>		
	ADDITION	AL COMMENTS:	
Prescriber Signature:		Date:	
		Dutt,	

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intended recipient, please notify us immediately by faxing back to the originator.

×