

 Please cut along the dotted lines before submitting to a pharmacy.

Date Required: _____ Ship To: _____ Patient _____ MD Office _____ Other: _____

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name: _____	Prescriber Name: _____
Address: _____	Address: _____
City, State, Zip: _____	City, State, Zip: _____
Home Phone: _____	Phone: _____
Cell Phone: _____	Fax: _____
Date of Birth: _____ Gender: _____	DEA #: _____ NPI #: _____
Emergency Contact: _____ Phone: _____	Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card.)

Primary Insurance: _____ ID: _____ Group: _____

Secondary Insurance: _____ ID: _____ Group: _____

Prescription Card: _____ ID: _____ BIN: _____ PCN: _____

To better serve your patient and facilitate insurance authorization, please complete the pertinent sections:

PATIENT DIAGNOSIS/CLINICAL INFORMATION

ICD-10 Code: _____ TB/PPD Test: Positive Negative Date Read: _____

Diagnosis: _____ Weight: _____ kg lbs Height: _____ cm in %BSA: _____

Prior Medication: _____ Allergies: _____ NKDA

Length of Treatment: _____ Injection Training/Home Health RN visit is necessary. Yes No

Reason for Discontinuation: _____ Site of Care: Home MD Office Other: _____

PRESCRIPTION INFORMATION

Medication:	Dose/Strength:	Directions:	Quantity:	Refills:
Keytruda®	100mg/4mL vial	Infuse 200mg IV over at least 30 minutes every 3 weeks Infuse 400mg IV over at least 30 minutes every 6 weeks		
Imfinzi®	120mg/2.4mL vial 500mg/10mL vial	Patients >= 30kg: Infuse 10mg/kg IV every 2 weeks Patients >=30kg: Infuse 1500mg IV every 4 weeks Patients < 30kg: Infuse 10mg/kg IV every 2 weeks Patients < 30kg: Infuse 20mg/kg IV every 4 weeks		
Opdivo®	40mg/4mL vial 100mg/10mL vial 120mg/12mL vial 240mg/24mL vial	Infuse 240mg IV every 2 weeks Infuse 360mg IV every 3 weeks Infuse 480mg IV every 4 weeks Other: _____		
Opdivo Qvantig®	600mg/10,000 units vial	Inject 600mg/10,000 units SUBQ every 2 weeks Inject 600mg/10,000 units SUBQ every 4 weeks Inject 900mg/15,000 units SUBQ every 3 weeks		
Yervoy®	50mg/10mL vial 200mg/40mL vial	3 mg/kg IV every 3 weeks for 4 doses 3 mg/kg IV every 3 weeks for 4 doses, followed by 3 mg/kg IV every 12 weeks for up to 4 additional doses.		
Opdivo® + Yervoy®		Opdivo 3mg/kg IV followed by Yervoy 1mg/kg IV on the same day, every 3 weeks for 4 doses Opdivo 1mg/kg IV followed by Yervoy 3mg/kg IV on the same day, every 3 weeks for 4 doses Opdivo 240mg IV followed by Yervoy 1mg/kg IV on the same day every 3 weeks for 4 doses Opdivo 360mg IV every 3 weeks and Yervoy 1mg/kg IV every 6 weeks Opdivo 3mg/kg IV every 2 weeks and Yervoy 1mg/kg IV every 6 weeks		
Other:				

PREMEDICATION ORDERS/OTHER MEDICATIONS

Flush Protocol

NaCl 0.9% 5mL Heparin 10 units per mL 250mL 0.9% NaCl for hydration

NaCl 0.9% 10mL Heparin 100 units per mL Other: _____

Premedications & Other Medications

Infusion supplies as per protocol Acetaminophen _____ mg PO prior to infusion

Anaphylaxis Kit orders as per protocol Diphenhydramine _____ mg PO

By signing this form and using this pharmacy's services, you are authorizing this pharmacy to serve as your prior authorization designated agent in dealing with prescription and medical insurance companies.

Prescriber Signature: _____ Date: _____