

 Fax Referral To: 877-277-9155
 Phone: 877-828-3940
 referrals@infucarerx.com
 www.infucarerx.health

Date Required: _____		Ship To: _____	Patient _____	MD Office _____	Other: _____
PATIENT INFORMATION			PRESCRIBER INFORMATION		
Patient Name: _____			Prescriber Name: _____		
Address: _____			Address: _____		
City, State, Zip: _____			City, State, Zip: _____		
Home Phone: _____			Phone: _____		
Cell Phone: _____			Fax: _____		
Date of Birth: _____ Gender: _____			DEA #: _____ NPI #: _____		
Emergency Contact: _____ Phone: _____			Contact Person: _____		
INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card.)					
Primary Insurance: _____		ID: _____	Group: _____		
Secondary Insurance: _____		ID: _____	Group: _____		
Prescription Card: _____		ID: _____	BIN: _____		PCN: _____
To better serve your patient and facilitate insurance authorization, please complete the pertinent sections:					
DIAGNOSIS		MEDICAL HISTORY			
E05.00 Thyrotoxicosis with diffuse goiter without thyrotoxic crisis or storm (hyperthyroidism)		Patient Weight: _____ kg lbs Height: _____ cm in			
Other: _____		Allergies: _____			
		Line Access: Peripheral Port			
		Delivery Method: Infusion Pump Other: _____			
		Nursing Coordination:			
		Pharmacy to coordinate home health nursing visit as necessary: Yes No			
		Home health nursing coordination not necessary. Reason:			
		MD office to administer to patient			
		Home health nursing already coordinated			
		(If applicable) Is the patient pregnant or planning to be pregnant? Yes No			
		Is the patient diabetic? Yes No			
		Does the patient have pre-existing or long term hearing loss? Yes No			
		Lab Orders: _____			
PRESCRIPTION INFORMATION					
Tepezza® (Teprotumumab-trbw) Prescription:				Quantity:	Refills:
Initiate services beginning with Dose No. _____ as indicated below:					
Dose 1: Infuse 10 mg/kg IV over 90 minutes, then 3 weeks later...					
Dose 2: Infuse 20 mg/kg IV over 90 minutes, then 3 weeks later...					
Dose 3 through 8: Infuse 20 mg/kg IV over 60 to 90 minutes (as tolerated by patient) every 3 weeks x 6 doses.					
PREMEDICATION ORDERS/OTHER MEDICATIONS					
Flush Protocol		Implanted Port:			
Peripheral:		NaCl 0.9% 5 to 10mL pre-/post-use and			
NaCl 0.9% 5mL		10 to 20mL pre-/post-lab draw			
NaCl 0.9% 10mL		Heparin (100 unit/mL) 3 to 5 mL post-use			
		For maintenance, heparin (100 unit/mL) 3 to 5mL every 24 hr if accessed or weekly to monthly if not accessed			
Premedications & Other Medications		Acetaminophen 650 mg PO prior to infusion			
Infusion supplies as per protocol		Diphenhydramine 25 mg PO			
Anaphylaxis Kit orders as per protocol		Other: _____			
ADDITIONAL COMMENTS:					

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intended recipient, please notify us immediately by faxing back to the originator.

Revised 31 August 2026